XV International AIDS Conference

Pathogenic Mechanisms of HIV Disease: The Role of Viral Replication and Immune Activation



Anthony S. Fauci, M.D.

National Institute of Allergy and Infectious
Diseases

National Institutes of Health Bethesda, Maryland, USA July 15, 2004





Viral Pathogenesis


Pathogenesis of HIV Disease




Immune Activation Drives HIV Pathogenesis



HIV Infection Leads to a State of Generalized Immune Activation



The Impact of ART on HIV-Associated Immune Activation



Reversal of Aberrant Immune Activation

Decreased T cell turnover


Increased T cell survival

Expansion of the CD4⁺ T cell pool

An immediate reversal in activation-induced immunodeficiency


- Spontaneous resolution of opportunistic illnesses
- Immune reactivation syndromes


Changes in CD4⁺ T Cell Production Following Initiation of Antiretroviral Therapy




Impact of Antigen and HIV Infection on the CD4⁺ T Cell Pool


Thymus








Impact of Viral Replication and Viremia on Lymphocyte Subsets in HIV-Infected Individuals



Impact of Viral Replication and Viremia on Lymphocyte Subsets in HIV-Infected Individuals





Presence of an Inducible HIV-1 Latent Reservoir During Highly Active Antiretroviral Therapy

Tae-Wook Chun, Lieven Stuyver, Stephanie B. Mizell, Linda A. Ehler, Jo Ann M. Mican, Michael Baseler, Alun L. Lloyd, Martin A. Nowak, and Anthony S. Fauci

PROCEEDINGS

OF THE

NATIONAL ACADEMY OF SCIENCES

OF THE UNITED STATES OF AMERICA

November 25, 1997 Volume 94 / Number 24

INCLUDES: FROM THE ACADEMY FEATURING GERMAN-AMERICAN FRONTIERS OF SCIENCE




Recovery of Replication-Competent HIV Despite Prolonged Suppression of Plasma Viremia

Joseph K. Wong, Marjan Hezareh, Huldrych F. Günthard, Diane V. Havlir, Caroline C. Ignacio, Celsa A. Spina, Douglas D. Richman


Identification of a Reservoir for HIV-1 in Patients on Highly Active Antiretroviral Therapy

Diana Finzi, Monika Hermankova, Theodore Pierson, Lucy M. Carruth, Christopher Buck, Richard E. Chaisson, Thomas C. Quinn, Karen Chadwick, Joseph Margolick, Ronald Brookmeyer, Joel Gallant, Martin Markowitz, David D. Ho, Douglas D. Richman, Robert F. Siliciano

Viral Relapse Following Discontinuation of ART



Importance of Viremia in Replenishment of the HIV Latent Reservoir



Effect of interleukin-2 on the pool of latently infected, resting CD4+ T cells in HIV-1-infected patients receiving highly active anti-retroviral therapy


Tae-Wook Chun, Richard T. Davey Jr, Delphine Engel, H. Clifford Lane, Anthony S. Fauci

Answers to the atypical measles puzzle

Autoimmune disease in mice lacking serum amyloid protein

Extracellular matrix proteins protect cancer cells

HAART, IL-2 and the eradication of HIV-1



New England Journal of Medicine

Established in 1812 as THE NEW ENGLAND JOURNAL OF MEDICINE AND SURGERY

VOLUME 340 MAY 27, 1999 NUMBER

Quantifying Residual HIV-1 Replication in Patients Receiving Combination Antiretroviral Therapy

Linqi Zhang, Bharat Ramratnam, Klara Tenner-Racz, Yuxian He, Mika Vesanen, Sharon Lewin, Andrew Talal, Paul Racz, Alan S. Perelson, Bette T. Korber, Martin Markowitz, and David D. Ho



Long-Term Follow-Up Studies Confirm the Stability of the Latent Reservoir for HIV-1 in Resting CD4⁺ T Cells

Janet D. Siliciano, Joleen Kajdas, Diana Finzi, Thomas C. Quinn, Karen Chadwick, Joseph B. Margolick, Colin Kovacs, Stephen J. Gange and Robert F. Siliciano

all are in jour liberes

PNAS

Proceedings of the National Academy of Sciences of the United States of America

September 28, 1999 | vol. 96 | no. 20 | pp. 10945–11688 | www.pnas.org

Perspective


Latent Reservoirs of HIV: Obstacles to the Eradication of Virus

Tae-Wook Chun and Anthony S. Fauci


The Resting CD4⁺ T Cell Reservoir of **HIV** in Viremic Versus Aviremic Individuals

Morphology of Resting CD4⁺ T Cells

Aviremic Patient




Viremic Patient




Resting CD4⁺ T Cell Reservoir for HIV in Viremic versus Aviremic Individuals is Phenotypically Identical

Patient 4 (Viremic)



Levels of Cell-Free HIV Virions Released by Latently Infected, Resting CD4⁺ T Cells from Viremic and Aviremic Patients



DNA Microarray Analysis of Resting CD4+ T Cells from Aviremic and Viremic HIV-Infected Patients

Isolation of RNA from resting CD4⁺ T cells



Synthesis and hybridization of cRNA onto Affymetrix Human Genome U95A Oligonucleotide Array (12,600 genes)




Analysis of data using Significant Analysis of Microarrays algorithm



Output of data using descriptive categories of genes using literature-mining algorithm


DNA Microarray Analysis of Resting CD4⁺ T Cells from Aviremic and Viremic Patients and Normal Donors



Conclusions

- There is a fundamental difference in the resting CD4⁺ T cell reservoir in viremic versus aviremic patients. In the former, true latency likely does not exist as cells are continually poised to express virus. In the latter, much greater stability exists and this may represent a truly latent reservoir of virus.
- Active viral replication, as manifested by detectable plasma viremia, has a significant impact on the physiologic state of resting CD4⁺ T cells in infected viremic patients, and in turn, allows release of cellfree HIV-1 without exogenous activation stimuli.

Impact of Viral Replication and Viremia on Lymphocyte Subsets in HIV-Infected Individuals



New England Journal of Medicine

Established in 1812 as THE NEW ENGLAND JOURNAL OF MEDICINE AND SURGERY

VOLUME 309 August 25, 1983 NUMBER 8

Abnormalities of B-Cell Activation and Immunoregulation in Patients with the Acquired Immunodeficiency Syndrome

H.C. Lane, H. Masur, L.C. Edgar, G. Whalen, A.H. Rook, and A.S. Fauci

PNAS

Proceedings of the National Academy of Sciences of the United States of America


August 28, 2001 | vol 98 | no. 18

HIV-1 Induces Phenotypic and Functional Perturbations of B Cells in Chronically Infected Individuals

Susan Moir, Angela Malaspina, Kisani M. Ogwaro, Eileen T. Donoghue, Claire W. Hallahan, Linda A. Ehler, Shuying Liu, Joseph Adelsberger, Réjean Lapointe, Patrick Hwu, Michael Baseler, Jan M. Orenstein, Tae-Wook Chun, Jo Ann M. Mican, and Anthony S. Fauci

Previous Findings Indicate That HIV-Mediated Hyper-Activation Induces Terminal Differentiation of B Cells

- Loss of CD21 expression
- Reduced proliferation
- Increased immunoglobulin secretion
- Changes in morphology



HIV-negative


HIV-viremic CD21+

HIV-viremic CD21-


Determination of B Cell Gene Expression in Viremic versus **Aviremic HIV-Infected** versus HIV-Negative Individuals

DNA Microarray Analysis: Genes Upregulated in B Cells of HIV-Viremic Patients

- Major categories of genes upregulated:
 - Interferon-stimulated genes
 - Genes associated with B cell terminal differentiation




Microarray Sub-analysis: Genes of Cell Surface Receptors Associated with Interferon Induction and B Cell Terminal Differentiation




Increased Susceptibility of B Cells of HIV-Viremic Patients to Fas-Mediated Apoptosis




B Cells Treated With FasL



Direct Correlation Between Level of Fas Expression on B Cells and Susceptibility of B Cells to Fas-Mediated Apoptosis




Direct Correlation Between HIV Plasma Viremia and Susceptibility of B Cells to Fas-Mediated Apoptosis




Summary

- A large number (~42) of B cell genes are upregulated in HIV-viremic vs HIV-aviremic and HIV-negative individuals; 75% are interferonstimulated or associated with B cell terminal differentiation.
- Changes in expression of TNFSF receptors such as CD95 (Fas) are associated with increased B cell death by apoptosis.

Impact of Viral Replication and Viremia on Lymphocyte Subsets in HIV-Infected Individuals



Function of NK Cells




2 Major Types of NK Cell Receptors

- Activating
- Inhibitory (Dominant)

NK Cell Activating Receptors

NK Cell Natural Cytotoxicity Receptors



Receptors NKp46 NKp30 NKp44 NKG2D

2B4 NTB-A DNAM-1 NKp80

(Signaling polypeptides)

 (\mathcal{G}_{γ})


 (\mathcal{Y}_{γ})

(DAP12)


(DAP10)

(LAT)


NK Cell Inhibitory Receptors



NK Cell Function




NK Cell Function






Natural Killer Cells in HIV-1 Infection: Dichotomous Effects of Viremia on Inhibitory and Activating Receptors and Their Functional Correlates

Domenico Mavilio et al




- HIV viremic patients manifest:
 - Decrease in activating receptors (NCRs)
 - Normal or increase in inhibitory receptors


Enrichment of CD56⁻/CD16⁺ NK Cells in HIV-1 Viremic Patients




The CD56⁻/CD16⁺ NK Cell Subset is Enriched in HIV-Viremic Patients



Inhibitory NK Receptors are Either Conserved or Increased in CD56⁻ NK Cells of HIV-Viremic Patients




Activating NK Receptors are Decreased in CD56⁻ NK Cells of HIV-Viremic Patients




Effect of HIV Viremia on CD56-NK Cells

- Decreased production of perforin and granzyme
- Decreased secretion of TNF-α and interferon-γ

NK Cells from HIV Viremic Individuals Express Fas on their Surface at a Significantly Higher Level than do NK Cells from Normal and HIV-Aviremic Individuals



NK Cells from HIV-Viremic Individuals Undergo Increased Apoptosis upon Exposure to sFasL *in vitro*




Effect of HIV Viremia on NK Cell Phenotype and Function


- HIV-viremic patients manifest a dichotomy of expression of NK cell receptors: inhibitory receptors are maintained or increased while activating receptors are decreased.
- This dichotomy is reflected in decreased cytotoxic function as well as a decrease in production of perforin and granzyme and a decrease in secretion of TNF-α and IFN-γ.
- Viremia is associated with an enrichment of CD56⁻/CD16⁺ subset of NK cells. The defects in receptor phenotype and function are concentrated within this abnormal subset.
- NK cells from viremic patients express increased levels of Fas on their surface and the CD56^{dim}/CD16+ subset responds to sFasL with increased apoptosis.

Direct Effects of HIV Envelope on Immune Cell Function


Role of HIV Envelope in the Induction of Aberrant Signal Transduction




The Role of NFAT in HIV Life Cycle




NFAT Activation through HIV Envelope Requires Coordinate Signaling through CD4 and Coreceptor




Categories of Genes Differentially Modulated by R5 vs X4 Envelopes




Cluster 4 - "Protein Modification" 0.6 R₅ 0.5 X4 Ln Fold Change 0.4 0.3 0.2 0.1 0.0 -0.1 -0.2-0.3 -0.416 5 Time (hrs.)








Conclusions


- HIV envelope modulates PBMC gene expression that is associated with virus replication and the secretion of cytokines involved in immune activation.
- This induction of gene expression requires coordinate signaling through CD4 and the respective co-receptor.
- Envelopes derived from X4 versus R5 viruses can differentially induce the expression of genes associated with transcription factors, cell cycle, and protein modification.
- Thus, HIV envelope-induced cell signaling may play an important role in the propagation of virus replication and the state of aberrant immune activation.






































NIAID Laboratory of Immunoregulation



Collaborators

NIAID/NIH

H. Clifford Lane
Richard Davey
Marybeth Daucher
Gabriella Lombardo
JoAnn M. Mican
J. Shawn Justement
Richard Lempicki
Jun Yang
Marie A. Planta

University of Genova

Alessandro Moretta, M.D.

Human Genome Sciences, Inc

Oxana K. Pickeral Charles E. Birse

University of Toronto

Colin Kovacs
Mona Loutfy
Rowena Dimayuga